

Uncovering mildew resistance biomarkers in grapevine: an integrated omics approach

Marisa Maia^{1,2,3,#}, Joana Figueiredo^{1,2,3}, Andreia Figueiredo^{3,*}, Marta Sousa Silva^{1,2,*}

¹ Laboratório de FTICR e Espectrometria de Massa Estrutural, FCUL, Portugal;

² Centro de Química e Bioquímica, FCUL, Portugal; ³ Biosystems & Integrative Sciences Institute (BioISI), FCUL, Portugal;

mrrmaia@fc.ul.pt; *co-senior authors

State of the art


Grapevine (*Vitis vinifera* L.) is one of the most important and cultivated fruit plant in the world due to its food products mainly to the wine industry. The European Union is the world's leader in wine production, with almost half of the total vine-growing area. Portugal is the fifth wine producing country and the second with the highest collection of grapevine varieties [1].

The domesticated *V. vinifera* cultivars frequently used for wine production are highly susceptible to fungal diseases, which if not controlled, presents serious negative effects for the country's economy. Downy and Powdery mildews are the most destructive vineyard diseases, affecting all the green parts of the vine (leaves, shoots and bunches), causing yield reduction with significant production losses and financial consequences [2].

An alternative strategy to fungicide applications for disease prevention is the development of new grapevine varieties through breeding programs: combining the high degree of resistance of wild *Vitis* species (resistant to pathogens) with good berry quality for wine production of the domesticated *V. vinifera* (susceptible). Hence, to understand the innate resistance mechanism of *Vitis*, the metabolic characterization of grapevine leaves is of upmost importance, not only because plants contain a unique metabolome that change with the environment, development and upon fungal infections, but also to find resistance-associated biomarkers.

Major goal

Identification of grapevine metabolic and protein biomarkers associated with resistance or susceptibility to fungal diseases that can be used as molecular screening traits in grapevine breeding programs



Acknowledgments

This work was supported by projects UID/MULTI/00612/2013, PESt-OE/QUI/UI0612/2013, PESt-OE/BIA/UI4046/2014, by the investigator FCT program IF/00819/2015 and grant SFRH/BD/116900/2016 from Fundação para a Ciência e Tecnologia (Portugal). We also had support from the Portuguese Mass Spectrometry Network (LISBOA-01-0145-FEDER-022125) and the Project EU_FT-ICR_MS, funded by the European Union's Horizon 2020 research and innovation programme under grant agreement nr. 731077.

References

- [1] Organisation of Vine and Wine. 2017 OIV Statistical Report on World Vitiviniculture. 2017.
- [2] Gessler C, Pertot I, Perazzolli M. Plasmopara viticola: a review of knowledge on downy mildew of grapevine and effective disease management. Phytopathol Mediterr 2011;50:3–44. doi:10.14601/Phytopathol_Mediterr-9360.
- [3] Maia M, Monteiro F, Sebastiana M, Marques AP, Ferreira AEN, Freire AP, et al. Metabolite extraction for high-throughput FTICR-MS-based metabolomics of grapevine leaves. EuPA Open Proteomics 2016;12:4–9. doi:10.1016/j.euprot.2016.03.002.